
Pleasant Lake Aquifer

Benson and Pierce Counties

Aquifer At-a-Glance					
Area	12.0 square miles				
Aquifer Type	Unconfined Surficial				
Major Land Uses over Aquifer	Grassland/Pasture (50%)				
(percentage of aquifer area covered in 2017) ¹	Crops (28%)				
Depth to Water (2021)*	6-47 feet				
Total Unique Wells Sampled	39				
Wells Sampled in 2021	17				
Samples Collected in 2021	19				
Years Sampled	1996, 2001, 2006, 2011, 2016, 2021				


- *Depths to water may vary seasonally, year to year, and across the aquifer
- Aguifer materials consist of sands and gravels that were deposited by streams moving meltwater away from glaciers during the last ice age.2
- The aguifer ranges from 4-98 feet thick and averages about 43 feet thick.2
- Domestic, irrigation, and stock wells are installed in the aquifer.
- The cities of Leeds and Rugby draw water from the aquifer.
- In North Dakota, permits are required to withdraw large quantities of groundwater. In 2020, 160 million gallons of permitted water were drawn from the aquifer; municipal use consumed the largest quantity of water. For more information on water use and permits, contact the North Dakota Department of Water Resources (dwr.nd.gov).





Pleasant Lake Aquifer Pleasant Lake Aquifer Major Surficial Aquifers

2020 Pleasant Lake aquifer permitted water use (from North Dakota Department of Water Resources (dwr.nd.gov)) \



About the Agricultural Ambient Groundwater Monitoring Program

- The North Dakota Department of Environmental Quality monitors a network of wells in approximately 50 surficial aguifers that are at elevated risk of agricultural contamination.
- Aquifers are sampled on a 5-year rotation.
- Monitoring began in 1992.
- The vast majority of these aquifers are located in central and eastern North Dakota.
- Water is tested for 21 general chemistry parameters, eight trace metals, and 64 pesticides.

References

US Department of Agriculture, 2017, National Agricultural Statistics Service Cropland Data Layer. Randich, P.G., 1977, Ground-Water Resources of Benson and Pierce Counties, North Dakota, North Dakota State Water Commission County Ground-Water Studies 18-Part 3, North Dakota Geological Survey Bulletin 59.


Water Chemistry

Is Aquifer Water High in...?


	Analyte	Result	2021 Median Concentration	Potential Effects
	Arsenic	Locally	0.008 mg/L	Skin or circulatory system damage, increased cancer risk
r	Iron	YES	3.79 mg/L	Metallic taste/odor, discoloration of surfaces
	Manganese	YES	0.37 mg/L	
?	Sodium	NO	15.1 mg/L	Taste, people with certain health conditions may need to limit intake
	Sulfate	NO	32.4 mg/L	Taste/odor, laxative effect for people not used to the water
	The state of the s			

For more information about Maximum Contaminant Levels (MCLs), health effects, and treatment options for these contaminants and more, see the NDDEQ's fact sheets (deq.nd.gov/wq/1_Groundwater) or visit the US EPA website (epa.gov/ground-water-and-drinking-water).


Dominant Water Type	Water Hardness
Calcium-Bicarbonate	Very Hard







Pesticides



State Pesticide Management Plan

Agricultural Groundwater Monitoring Program aquifers are monitored as a part of the State Pesticide Management Plan. A Prevention Action Level (PAL) threshold of 25% of the pesticide's Maximum Contaminant Level (MCL)* or Health Advisory Level (HAL) is used to identify whether action is needed to prevent further contamination.

Prevention Action Level Exceedances	None
MCL or HAL Exceedances	None

Number of Unique Wells with
Pesticide Detections since 1996

5 of 39 Total Wells

2021 Pesticide Detections				
Picloram	1 Well	Herbicide applied to crops and roads/rights-of-way		